



Review Article

Open Access

IMPROVEMENT OF WHEAT PRODUCTION THROUGH GENETIC AND ENVIRONMENTAL INTERACTIONS

ABBAS A¹, REHMAN AU¹, ALI T², SAMI A¹, HAIDER MZ¹

¹Department of Plant Breeding and Genetics, Faculty of Agricultural Sciences, University of the Punjab, P.O BOX. 54590, Lahore, Pakistan

²Department of Plant Breeding and Genetics University of Agriculture, Faisalabad, Pakistan

*Correspondence author email address: ali.bukhari91112@gmail.com

(Received, 30th September 2022, Revised 26th May 2023, Published 27th May 2023)

Abstract The effect of genetics and environmental factors on wheat production and the necessity for researchers to find genetic sources that can survive changing environmental variables to maximize crop yield. According to previous studies, despite the genetic history of the cultivated variety, drought, high salt, heat, infections, and microbes affect wheat development and yield. Climate change can provide unexpected weather that harms crop output, threatening agricultural sustainability. Researchers are exploring genotyping, phenological traits, GWAS, and QTL mapping to boost crop output and fulfill population needs. Breeding and genetic modification are essential for wheat production due to a growing population, changing climate, and limited planted areas. The article also mentioned weighing 1000 grains or kg per hectare to measure yield.

Keywords: Environmental factor, Microbes, Sustainability, GWAS, QTL mapping

Introduction

Triticum aestivum and *Triticum durum* are two types of wheat that are essential protein sources and calories for millions of people worldwide (Arzani & Ashraf, 2017). It provides 18–20% of the world's calories and is a mainstay crop, making a big difference in ensuring enough food for everyone (Rampa et al., 2020). The demand for wheat is likely to keep growing because the world's population and eating habits are constantly changing (Enghiad et al., 2017; Fróna et al., 2019). We currently grow more than 200 million hectares of wheat, the most grains of any crop (Stewart & Lal, 2018). To keep up with the growing demand for food and ensure enough for everyone, it is important to create high-yield wheat types that can handle harsh conditions (Liliane & Charles, 2020; Zörb et al., 2018). Asseng et al. (2017) have found several factors that affect output, such as the number of spikelet's per spike and the weight of 1000 kernels that help wheat grow well. So, to get the highest yields, breeding programs try to make wheat plants with these traits (Venske et al., 2019). However, global wheat yields have decreased since the 1990s, and climate change is still a threat. This means there is a need to quickly create wheat varieties that can handle abiotic stresses like drought, temperature changes, and soil salinization (Kumaraswamy & Shetty, 2016; Mariani & Ferrante, 2017; Sarwar et al., 2021).

To solve this problem, scientists have been looking into the different types of wheat and finding genotypes that can handle these stressful situations better and produce more crops (Beres, Hatfield, et al., 2020; Beres, Rahmani, et al., 2020). Molecular markers that can speed up the creation of wheat types that can handle stress are also being looked for in these studies (Raza et al., 2019; Younis et al., 2020). Making high-yielding and stress-resistant wheat types is key to ensuring enough food for everyone worldwide. As the world's population and climate change, we need to quickly create new types of wheat that can thrive in these circumstances. The growth of new wheat types can meet the growing need for food by looking into the genetic diversity of wheat and using molecular markers in breeding programs (Ali et al., 2013; Alipour et al., 2017; Ghafoor et al., 2020; Wani et al., 2022).

Different Techniques use for Crop Yield Improvement

According to experts, improving factors like genotype, surroundings, and phenological traits could raise yields by 43-62%. These factors include how the leaves look, how they react to the length of the days for blooming, how much photosynthesis can happen through the flag leaf area, how well the plant can handle drought, and how well the roots can take water (Iqra et al., 2020; Muqadas et al., 2020; Takeno, 2016). Many experts are working on different types of wheat to improve important traits like photosynthesis

and leaf gas exchange, as well as grain number and size. These optimized source and sink traits must work together well to increase food yields. Many characteristics, like the number of tillers per plant, the 1000-kernel weight, the time of flowering, the days until heading, and the length of the spike, have been looked at to see how they affect output ([Mathan et al., 2016](#); [Naseem et al., 2020](#); [Nawaz et al., 2020](#)). It is important to know that the number of spikes on a plant is linked to the number of tillers, affecting the grain yield of hexaploid wheat. Furthermore, breeding programs have shown that wheat types with higher grain weight have more chlorophyll pigment, affecting grain weight in direct and indirect ways. Multiple wheat genotypes had grain weights ranging from 20 to 50 g per 1000 kernels in one study ([Ali, 2017](#); [Sarwar et al., 2022](#); [Shafique et al., 2020](#)). This was due to grain filling and cell growth through water absorption. Genotypes with better root growth, biomass, and water uptake have higher grain weight and better grain filling. Scientists have also looked at other factors that affect 1000-kernel weight, such as the weight of the carpel at anthesis, the date of anthesis for each flower, and the length of time the grains are filled ([Wu et al., 2018](#)).

Genotype Effects on Wheat Yield

Wheat is a significant food item worldwide, a hexaploid plant with the genetic code $2n=6x=42$ and a primarily studied family name (Poaceae). The plant is polyploid, which means it has three different genomes, each with two sets of basic chromosomes ($2n=2x=14$). This species has three genomes: the A genome comes from *Triticum urartu*, the B genome comes from *Aegilops speltoides*, and the D genome comes from *A. Tauschii*. This complicates and is genetically varied ([Mirzaghaderi & Mason, 2019](#)). Different types of wheat have different genetic variations that affect their yield. Plant height, the number of tillers and spikelet's, grain weight, and other factors affect wheat's total productivity ([Ali et al., 2016](#)). [Li et al. \(2016\)](#) found that the number of spikelets per spike varies between kinds of wheat. The number of tillers per plant can also affect the number of spikelet's and the weight of the grain. Earlier research has shown that several farming and physiological traits are genetically variable. These include plant height, tiller and kernel number, days to maturity, and blooming ([Voss-Fels et al., 2019](#)). Genetic prediction works better for smaller groups of people than QTL-based prediction ([Norman et al., 2017](#)). Genetic and morphological studies of wheat spike and kernel traits have shown geographical patterns and long-term trends. These studies are mostly about spikelet fertility and kernel/spikelet number, determining the final grain yield. Genome prediction has shown that these qualities are passed down quantitatively and are affected by small-effect QTLs ([Moore et al., 2017](#)).

Yield Improvement using GWAS and QTL Mapping

It was hopeful at first that linkage mapping and GWAS would significantly improve grain yield because of how complicated yield is, both genetically and in terms of how it interacts with biotic and abiotic factors in the field. But, despite attempts, this hasn't happened. QTL mapping is a powerful tool for finding specific genome regions and learning about the physiological and genetic processes that control important characteristics ([Bhanu et al., 2016](#)). This method works well for quality traits that don't depend on many genes. Additionally, Marker-Assisted Selection using closely related markers or the causing mutation can help choose phenotypes more effectively ([Visscher et al., 2021](#)). However, it has been hard to find important QTLs and useful markers for yield. Some studies found that the allele from one parent increased grain output, while others found that the allele from the other parent did a better job. We don't fully understand the natural factors that cause this variation yet. Some things that could change are the number of seeds planted, how they are watered, and the stress caused by heat and drought. It's hard to find yield QTLs and useful selection markers, as this shows ([Platten et al., 2019](#); [Yang et al., 2020](#)).

Focusing on key sites within physiological systems instead of individual QTLs is better than keeping yields stable across various environmental factors ([Dwivedi et al., 2020](#)). In places with many terminal droughts, for example, wheat and barley plants may do better if their roots are small to reach deep water in the ground ([Daryanto et al., 2017](#)). On the other hand, breeding rice to be more resistant to drought and better at establishing itself in flooded fields could mean making rice plants more resistant to senescence and anaerobic germination ([Kumar et al., 2017](#)). These particular traits are more important in specific settings than in terms of progress as a whole. Many links have been found between adaptive traits and field yield ([Furbank et al., 2019](#)). So, we must look at the whole picture to determine how genetics affects yield-related traits. This means using methods that combine genetics, physiology, and metabolism to figure out how things work ([Sallam et al., 2019](#)). Field testing in natural farming conditions is significant, and using a mix of methods like linkage mapping and GWAS has been shown to work best. It's now easier to do this with the help of freely available datasets and tools ([Shadrin et al., 2021](#)). For advanced methods like GWAS and forward and reverse genetic approaches ([Kondić-Špika et al., 2023](#); [Walli et al., 2022](#)), high-throughput phenotyping is necessary.

Influence of Biotic and Abiotic Stress Factors on Wheat Yield

Healthy Environment Soil Salinization Management

Abiotic stress affecting field crops is mainly caused by salt, with drought being the other major. Indicators like [Shankar and Prasad \(2023\)](#) say this stress can significantly lower food yield, growth, and production. The physical and chemical qualities of the

soil, as well as the ecosystem's balance, are also greatly affected. Soil erosion, low food yields, and lower economic returns are some problems that can happen because of salinity stress ([El Sabagh et al., 2020](#)).

Plants' anatomy, physiology, and biochemistry are all damaged by salinity stress, which makes it harder for seeds to sprout, grow, and take in water and nutrients ([Shahid et al., 2020](#)). Ion poisoning, a lack of important elements like N, Ca, K, P, Fe, and Zn, osmotic stress, and less water intake are all caused by it. Beyond that, plants can also be hurt by too much sodium, chlorine, and boron ([Chakraborty et al., 2018](#)). Photosynthesis is directly affected as the leaf area, chlorophyll content, and stomatal opening all drop ([Chakraborty et al., 2018](#); [Hu et al., 2020](#)). According to [Bhattacharjya et al. \(2018\)](#), salinity stress can also affect reproductive development. It can stop micro-sporogenesis and stamen filament lengthening, cause more tissue cell death, lead to ovule abortion, and slow embryo development. Plants' growth and output are greatly affected by salinity stress. In the worst cases, plants can die, and resources are lost. Two ways plants deal with salinity are by either getting rid of too many salts in their cells or being okay with them being there ([Acosta-Motos et al., 2017](#); [Dassanayake & Larkin, 2017](#)). Varying amounts of salt can hurt different foods. Wheat (*T. aestivum*), a grain that can handle salt, does best in mild climates, while rice (*Oryza sativa* L.), on the other hand, is susceptible to salt and can't grow in salty places ([Hasan et al., 2022](#)). Salt-tolerant crops like barley (*Hordeum vulgare*) can still die in high salt levels, but durum wheat is not as salt-tolerant as bread wheat ([Mwando, 2021](#)). Biochemical, physiological, and molecular levels all show that salt stress hurts plant growth and development as a whole ([Dogan, 2020](#); [Isayenkov & Maathuis, 2019](#)).

Effect of Drought Stress

The main reason for drought in agriculture is insufficient water in the root zone, resulting in less production. Drought has many effects that are different in different places. To be able to survive drought, plants need adaptation, which includes the ability to avoid drought and the ability to handle being dry. However, this tolerance differs from salt tolerance ([Abobatta, 2020](#); [Asif et al., 2020](#); [Ghafoor et al., 2020](#); [Ilyas et al., 2021](#); [Zubair et al., 2016](#)). Controlling how much the plant absorbs and moves the salt is very important for salt tolerance. Depending on the climate, drought can have different effects on crop production. For example, traits or genes that increase yield in severe drought may not work in moderate drought or may even have adverse effects when there is enough water. For this reason, we need to improve plants' ability to handle drought in certain areas. Plants have many ways to fight off drought, such as controlling the behavior of their stomata, the balance of osmotic pressure, the activity of antioxidants, hormone signaling, miRNA expression,

light protection, and metabolic pathways ([dos Santos et al., 2022](#); [Kaur et al., 2021](#)). Dry conditions change gene expression and metabolite output, meaning many genes and pathways are involved in this regulatory network. Photosynthesis is crucial during the blooming phase because it creates a good sink potential for later higher output ([Ahmad et al., 2021](#); [Ali et al., 2016](#); [Soltani et al., 2019](#)). Many things affect grains, including carbon and nitrogen availability, 10 to 15 days before anthesis. As weather conditions worsen, a lack of water from insufficient rain and groundwater loss are major threats to crop production and food security. To feed everyone, crop production needs to improve, even if there isn't a lot of water available ([McAfee, 2019](#)). Because of climate change, droughts, and storms will happen more often, making it even harder to grow food.

Effect of temperature

Stress from heat harms wheat yield. Asseng et al. (2015) say that average world wheat yields will drop by 4 to 6 percent for every 1°C rise in global mean temperature ([Zhao et al., 2017](#)). Heat can affect growth and grain filling depending on when and how strong the stress is ([Sehgal et al., 2018](#)). Heat stress can make flag leaf photosynthesis less effective, which makes it harder for assimilates to build up and move around. This can lead to a significant drop in grain yield and quality ([Feller, 2016](#)).

Effect of Phyto-pathogens

The agricultural sector faces many problems as the world's population grows quickly. Plant diseases can hurt crop output, leading to lower yields, especially for crops used for biofuel and fiber. This can be very bad or long-lasting ([Antar et al., 2021](#)). For example, the top six food crops lose an average of 45% of their production. Postharvest disease can also be very bad, especially when it's hard to get to markets, the infrastructure isn't good, and the supply chain isn't working well. Some of these diseases can also make poisons that harm people's health. Experts think that changes in the climate, like global warming, can make plant germs spread faster and cause more damage ([Bhadra et al., 2022](#)). According to [Petronaitis et al. \(2021\)](#), these pathogens cause crop losses of 20 to 30 percent annually. This costs a lot of money and affects food security locally, nationally, and internationally. The direct effect of Phytopathogens on grain yields is responsible for 25–45% of all farm production worldwide ([Petronaitis et al., 2021](#)).

Effect of Photosynthesis

Photosynthesis has become more and more popular as a way to improve crop yield over the years. Photosynthetic models have helped researchers learn about different parts of photosynthetic leaf gas exchange. Some of these studies are by ([Wu et al., 2019](#)) created a diurnal canopy photosynthesis stomatal conductance model that shows how leaf photosynthesis affects crop output in an Australian cropping region in the spring and summer, when

water is limited and when water is not limited (Wu et al., 2019).

Conclusion

The last few decades have seen a lot of progress in our knowledge of the genes and processes that control important traits that increase crop yield, like the amount and size of grains, the number and function of stomata, the structure of leaves, and the storage of carbohydrates. However, regarding grain output and wheat quality, how genotypes and environmental conditions interact often leaves people guessing what will happen. This is a big problem for systems that look at inbreeding and genotypes. Soil quality, nitrogen supply, rainfall, and temperatures during ripening are critical factors that affect how well different genotypes do. To make a single gene change in wheat work, you need to simultaneously target all three copies of the gene using methods like gene recognition and editing. GWAS and QTL mapping are both thorough ways to increase crop yields, but they haven't significantly impacted total crop productivity yet. Gene editing has a lot of promise, especially regarding decreasing negative yield regulators. Researchers can speed up the creation of drought- and stress-resistant crop types by finding and using new molecular markers. It's important to balance vegetative and root growth for significant output gains and make the most of the carbon allocation processes during grain set and filling. To do this, you need to know how genes and systems work together in the field. Basic study is still ongoing, and gene editing could be helpful. This gives us reason to be hopeful about getting significant yield gains in the future.

References

- Abobatta, W. F. (2020). Plant responses and tolerance to combined salt and drought stress. *Salt and Drought Stress Tolerance in Plants: Signaling Networks and Adaptive Mechanisms*, 17-52.
- Acosta-Motos, J. R., Ortuño, M. F., Bernal-Vicente, A., Diaz-Vivancos, P., Sanchez-Blanco, M. J., & Hernandez, J. A. (2017). Plant responses to salt stress: adaptive mechanisms. *Agronomy*, 7(1), 18.
- Ahmad, M., Ali, Q., Hafeez, M., & Malik, A. (2021). Improvement for biotic and abiotic stress tolerance in crop plants. *Biological and Clinical Sciences Research Journal*, 2021(1).
- Ali, M. J. (2017). Investigation of yield, yield components and primary quality characteristics of some bread wheat (*Triticum aestivum* L.) genotypes.
- Ali, Q., Ahsan, M., Ali, F., Aslam, M., Khan, N. H., Munzoor, M., Mustafa, H. S. B., & Muhammad, S. (2013). Heritability, heterosis and heterobeltiosis studies for morphological traits of maize (*Zea mays* L.) seedlings. *Advancements in Life Sciences*, 1(1).
- Ali, Q., Ahsan, M., Kanwal, N., Ali, F., Ali, A., Ahmed, W., Ishfaq, M., & Saleem, M. (2016). Screening for drought tolerance: comparison of maize hybrids under water deficit condition. *Advancements in Life Sciences*, 3(2), 51-58.
- Alipour, H., Bihamta, M. R., Mohammadi, V., Peyghambari, S. A., Bai, G., & Zhang, G. (2017). Genotyping-by-sequencing (GBS) revealed molecular genetic diversity of Iranian wheat landraces and cultivars. *Frontiers in Plant Science*, 8, 1293.
- Antar, M., Lyu, D., Nazari, M., Shah, A., Zhou, X., & Smith, D. L. (2021). Biomass for a sustainable bioeconomy: An overview of world biomass production and utilization. *Renewable and Sustainable Energy Reviews*, 139, 110691.
- Arzani, A., & Ashraf, M. (2017). Cultivated ancient wheats (*Triticum* spp.): A potential source of health-beneficial food products. *Comprehensive Reviews in Food Science and Food Safety*, 16(3), 477-488.
- Asif, S., Ali, Q., & Malik, A. (2020). Evaluation of salt and heavy metal stress for seedling traits in wheat. *Biological and Clinical Sciences Research Journal*, 2020(1).
- Asseng, S., Kassie, B. T., Labra, M. H., Amador, C., & Calderini, D. F. (2017). Simulating the impact of source-sink manipulations in wheat. *Field Crops Research*, 202, 47-56.
- Beres, B. L., Hatfield, J. L., Kirkegaard, J. A., Eigenbrode, S. D., Pan, W. L., Lollato, R. P., Hunt, J. R., Strydhorst, S., Porcher, K., & Lyon, D. (2020). Toward a better understanding of genotype× environment× management interactions—a global wheat initiative agronomic research strategy. *Frontiers in Plant Science*, 11, 828.
- Beres, B. L., Rahmani, E., Clarke, J. M., Grassini, P., Pozniak, C. J., Geddes, C. M., Porcher, K. D., May, W. E., & Ransom, J. K. (2020). A systematic review of durum wheat: Enhancing production systems by exploring genotype, environment, and management (G× E× M) synergies. *Frontiers in Plant Science*, 11, 568657.
- Bhadra, P., Maitra, S., Shankar, T., Hossain, A., Praharaj, S., & Aftab, T. (2022). Climate change impact on plants: Plant responses and adaptations. In *Plant perspectives to global climate changes* (pp. 1-24). Elsevier.
- Bhanu, A., Singh, M., Srivastava, K., & Hemantaranjan, A. (2016). Molecular mapping and breeding of physiological traits. *Adv. Plants Agric. Res.*, 3(00120), 10.15406.
- Bhattacharjya, S., Bhaduri, D., & Sahu, A. (2018). Arbuscular mycorrhizal fungi: a potential tool for enhancing crop productivity in salt affected soil. *International Journal of Agriculture, Environment and Biotechnology*, 11(6), 871-880.
- Chakraborty, K., Basak, N., Bhaduri, D., Ray, S., Vijayan, J., Chattopadhyay, K., & Sarkar, R. K. (2018). Ionic basis of salt tolerance in plants:

- nutrient homeostasis and oxidative stress tolerance. *Plant nutrients and abiotic stress tolerance*, 325-362.
- Daryanto, S., Wang, L., & Jacinthe, P.-A. (2017). Global synthesis of drought effects on cereal, legume, tuber and root crops production: A review. *Agricultural Water Management*, 179, 18-33.
- Dassanayake, M., & Larkin, J. C. (2017). Making plants break a sweat: the structure, function, and evolution of plant salt glands. *Frontiers in Plant Science*, 8, 406.
- Dogan, M. (2020). Effect of salt stress on in vitro organogenesis from nodal explant of *Limnophila aromatica* (Lamk.) Merr. and *Bacopa monnieri* (L.) Wettst. and their physiological and biochemical responses. *Physiology and molecular biology of plants*, 26, 803-816.
- dos Santos, T. B., Ribas, A. F., de Souza, S. G. H., Budzinski, I. G. F., & Domingues, D. S. (2022). Physiological responses to drought, salinity, and heat stress in plants: a review. *Stresses*, 2(1), 113-135.
- Dwivedi, S. L., Stoddard, F. L., & Ortiz, R. (2020). Genomic-based root plasticity to enhance abiotic stress adaptation and edible yield in grain crops. *Plant Science*, 295, 110365.
- El Sabagh, A., Hossain, A., Barutçular, C., Iqbal, M. A., Islam, M. S., Fahad, S., Sytar, O., Çiğ, F., Meena, R. S., & Erman, M. (2020). Consequences of salinity stress on the quality of crops and its mitigation strategies for sustainable crop production: an outlook of arid and semi-arid regions. *Environment, climate, plant and vegetation growth*, 503-533.
- Enghiad, A., Ufer, D., Countryman, A. M., & Thilmany, D. D. (2017). An overview of global wheat market fundamentals in an era of climate concerns. *International Journal of Agronomy*, 2017.
- Feller, U. (2016). Drought stress and carbon assimilation in a warming climate: Reversible and irreversible impacts. *Journal of Plant Physiology*, 203, 84-94.
- Fróna, D., Szenderák, J., & Harangi-Rákos, M. (2019). The challenge of feeding the world. *Sustainability*, 11(20), 5816.
- Furbank, R. T., Jimenez-Berni, J. A., George-Jaeggli, B., Potgieter, A. B., & Deery, D. M. (2019). Field crop phenomics: enabling breeding for radiation use efficiency and biomass in cereal crops. *New Phytologist*, 223(4), 1714-1727.
- Ghafoor, M., Ali, Q., & Malik, A. (2020). Effects of salicylic acid priming for salt stress tolerance in wheat. *Biological and Clinical Sciences Research Journal*, 2020(1).
- Hasan, M. N., Bhuiyan, F. H., Hoque, H., Jewel, N. A., Ashrafuzzaman, M., & Prodhana, S. H. (2022). Ectopic expression of *Vigna radiata*'s vacuolar Na⁺/H⁺ antiporter gene (VrNHX1) in indica rice (*Oryza sativa* L.). *Biotechnology Reports*, 35, e00740.
- Hu, W., Lu, Z., Meng, F., Li, X., Cong, R., Ren, T., Sharkey, T. D., & Lu, J. (2020). The reduction in leaf area precedes that in photosynthesis under potassium deficiency: the importance of leaf anatomy. *New Phytologist*, 227(6), 1749-1763.
- Ilyas, M., Nisar, M., Khan, N., Hazrat, A., Khan, A. H., Hayat, K., Fahad, S., Khan, A., & Ullah, A. (2021). Drought tolerance strategies in plants: a mechanistic approach. *Journal of Plant Growth Regulation*, 40, 926-944.
- Iqra, L., Rashid, M. S., Ali, Q., Latif, I., & Mailk, A. (2020). Evaluation for Na⁺/K⁺ ratio under salt stress condition in wheat. *Life Sci J*, 17(7), 43-47.
- Isayenkov, S. V., & Maathuis, F. J. (2019). Plant salinity stress: many unanswered questions remain. *Frontiers in Plant Science*, 10, 80.
- Kaur, H., Kohli, S. K., Khanna, K., & Bhardwaj, R. (2021). Scrutinizing the impact of water deficit in plants: Transcriptional regulation, signaling, photosynthetic efficacy, and management. *Physiologia Plantarum*, 172(2), 935-962.
- Kondić-Špika, A., Trkulja, D., Brbaklić, L., Mikić, S., Glogovac, S., Johansson, E., Alemu, A., Chawade, A., Rahmatov, M., & Ibba, M. I. (2023). Marker-assisted selection for the improvement of cereals and pseudocereals. In *Developing Sustainable and Health Promoting Cereals and Pseudocereals* (pp. 253-283). Elsevier.
- Kumar, A., Sandhu, N., Yadav, S., Pradhan, S. K., Anandan, A., Pandit, E., Mahender, A., & Ram, T. (2017). Rice varietal development to meet future challenges. In *The future rice strategy for India* (pp. 161-220). Elsevier.
- Kumaraswamy, S., & Shetty, P. (2016). Critical abiotic factors affecting implementation of technological innovations in rice and wheat production: a review. *Agricultural reviews*, 37(4).
- Li, Y., Cui, Z., Ni, Y., Zheng, M., Yang, D., Jin, M., Chen, J., Wang, Z., & Yin, Y. (2016). Plant density effect on grain number and weight of two winter wheat cultivars at different spikelet and grain positions. *PLoS one*, 11(5), e0155351.
- Liliane, T. N., & Charles, M. S. (2020). Factors affecting yield of crops. *Agronomy-climate change & food security*, 9.
- Mariani, L., & Ferrante, A. (2017). Agronomic management for enhancing plant tolerance to abiotic stresses—drought, salinity, hypoxia, and lodging. *Horticulturae*, 3(4), 52.
- Mathan, J., Bhattacharya, J., & Ranjan, A. (2016). Enhancing crop yield by optimizing plant developmental features. *Development*, 143(18), 3283-3294.

- McAfee, A. (2019). *More from less: The surprising story of how we learned to prosper using fewer resources—And what happens next*. Scribner.
- Mirzaghaderi, G., & Mason, A. S. (2019). Broadening the bread wheat D genome. *Theoretical and Applied Genetics*, 132, 1295-1307.
- Moore, J. K., Manmathan, H. K., Anderson, V. A., Poland, J. A., Morris, C. F., & Haley, S. D. (2017). Improving genomic prediction for pre-harvest sprouting tolerance in wheat by weighting large-effect quantitative trait loci. *Crop Science*, 57(3), 1315-1324.
- Muqadas, S., Ali, Q., & Malik, A. (2020). Genetic association among seedling traits of Zea mays under multiple stresses of salts, heavy metals and drought. *Biological and Clinical Sciences Research Journal*, 2020(1).
- Mwando, E. K. (2021). *The genetics of barley (Hordeum vulgare) salinity tolerance during germination and the instantaneous seedling endurance* [Murdoch University].
- Naseem, S., Ali, Q., & Malik, A. (2020). Evaluation of maize seedling traits under salt stress. *Biological and Clinical Sciences Research Journal*, 2020(1).
- Nawaz, A., Haseeb, A., Malik, H., Ali, Q., & Malik, A. (2020). Genetic association among morphological traits of Zea mays seedlings under salt stress. *Biological and Clinical Sciences Research Journal*, 2020(1).
- Norman, A., Taylor, J., Tanaka, E., Telfer, P., Edwards, J., Martinant, J.-P., & Kuchel, H. (2017). Increased genomic prediction accuracy in wheat breeding using a large Australian panel. *Theoretical and Applied Genetics*, 130, 2543-2555.
- Petronaitis, T., Simpfendorfer, S., & Hüberli, D. (2021). Importance of Fusarium spp. in wheat to food security: A global perspective. *Plant diseases and food security in the 21st century*, 127-159.
- Platten, J. D., Cobb, J. N., & Zantua, R. E. (2019). Criteria for evaluating molecular markers: Comprehensive quality metrics to improve marker-assisted selection. *PloS one*, 14(1), e0210529.
- Rampa, F., Lammers, E., Linnemann, A., Schoustra, S., & de Winter, D. (2020). *Pathways to improved food and nutrition security of the poor: The promise of African indigenous foods and technologies*.
- Raza, A., Mehmood, S. S., Shah, T., Zou, X., Yan, L., Zhang, X., & Khan, R. S. A. (2019). Applications of molecular markers to develop resistance against abiotic stresses in wheat. *Wheat Production in Changing Environments: Responses, Adaptation and Tolerance*, 393-420.
- Sallam, A., Alqudah, A. M., Dawood, M. F., Baenziger, P. S., & Börner, A. (2019). Drought stress tolerance in wheat and barley: advances in physiology, breeding and genetics research. *International journal of molecular sciences*, 20(13), 3137.
- Sarwar, M., Anjum, S., Alam, M. W., Ali, Q., Ayyub, C., Haider, M. S., Ashraf, M. I., & Mahboob, W. (2022). Triacanol regulates morphological traits and enzymatic activities of salinity affected hot pepper plants. *Scientific reports*, 12(1), 3736.
- Sarwar, M., Anjum, S., Ali, Q., Alam, M. W., Haider, M. S., & Mehboob, W. (2021). Triacanol modulates salt stress tolerance in cucumber by altering the physiological and biochemical status of plant cells. *Scientific reports*, 11(1), 24504.
- Sehgal, A., Sita, K., Siddique, K. H., Kumar, R., Bhogireddy, S., Varshney, R. K., HanumanthaRao, B., Nair, R. M., Prasad, P. V., & Nayyar, H. (2018). Drought or/and heat-stress effects on seed filling in food crops: impacts on functional biochemistry, seed yields, and nutritional quality. *Frontiers in Plant Science*, 9, 1705.
- Shadrin, A. A., Kaufmann, T., van der Meer, D., Palmer, C. E., Makowski, C., Loughnan, R., Jernigan, T. L., Seibert, T. M., Hagler, D. J., & Smeland, O. B. (2021). Vertex-wise multivariate genome-wide association study identifies 780 unique genetic loci associated with cortical morphology. *Neuroimage*, 244, 118603.
- Shafique, F., Ali, Q., & Malik, A. (2020). Effects of heavy metal toxicity on maize seedlings growth traits. *Biological and Clinical Sciences Research Journal*, 2020(1).
- Shahid, M. A., Sarkhosh, A., Khan, N., Balal, R. M., Ali, S., Rossi, L., Gómez, C., Mattson, N., Nasim, W., & Garcia-Sanchez, F. (2020). Insights into the physiological and biochemical impacts of salt stress on plant growth and development. *Agronomy*, 10(7), 938.
- Shankar, A., & Prasad, V. (2023). Potential of desiccation-tolerant plant growth-promoting rhizobacteria in growth augmentation of wheat (*Triticum aestivum* L.) under drought stress. *Frontiers in Microbiology*, 14, 1017167.
- Soltani, A., Weraduwege, S. M., Sharkey, T. D., & Lowry, D. B. (2019). Elevated temperatures cause loss of seed set in common bean (*Phaseolus vulgaris* L.) potentially through the disruption of source-sink relationships. *BMC genomics*, 20(1), 1-18.
- Stewart, B., & Lal, R. (2018). Increasing world average yields of cereal crops: It's all about water. *Advances in agronomy*, 151, 1-44.
- Takeno, K. (2016). Stress-induced flowering: the third category of flowering response. *Journal of Experimental Botany*, 67(17), 4925-4934.
- Venske, E., Dos Santos, R. S., Busanello, C., Gustafson, P., & Costa de Oliveira, A. (2019). Bread wheat: a role model for plant

- domestication and breeding. *Hereditas*, 156(1), 1-11.
- Vissscher, P., Van der Beek, S., & Haley, C. (2021). Marker assisted selection. In *Animal Breeding* (pp. 119-136). Routledge.
- Voss-Fels, K. P., Cooper, M., & Hayes, B. J. (2019). Accelerating crop genetic gains with genomic selection. *Theoretical and Applied Genetics*, 132, 669-686.
- Walli, M. H., Al-Jubouri, Z., Madumarov, M. M., Margaryta, M., & Aldibe, A. A. A. (2022). Genetic and environment diversity to improve wheat (*Triticum* spp.) productivity: A review. *Research on Crops*, 23(2).
- Wani, S. H., Khan, H., Riaz, A., Joshi, D. C., Hussain, W., Rana, M., Kumar, A., Athiyannan, N., Singh, D., & Ali, N. (2022). Genetic diversity for developing climate-resilient wheats to achieve food security goals. *Advances in agronomy*, 171, 255-303.
- Wu, A., Hammer, G. L., Doherty, A., von Caemmerer, S., & Farquhar, G. D. (2019). Quantifying impacts of enhancing photosynthesis on crop yield. *Nature plants*, 5(4), 380-388.
- Wu, X., Tang, Y., Li, C., & Wu, C. (2018). Characterization of the rate and duration of grain filling in wheat in southwestern China. *Plant Production Science*, 21(4), 358-369.
- Yang, L., Zhao, D., Meng, Z., Xu, K., Yan, J., Xia, X., Cao, S., Tian, Y., He, Z., & Zhang, Y. (2020). QTL mapping for grain yield-related traits in bread wheat via SNP-based selective genotyping. *Theoretical and Applied Genetics*, 133, 857-872.
- Younis, A., Ramzan, F., Ramzan, Y., Zulfiqar, F., Ahsan, M., & Lim, K. B. (2020). Molecular markers improve abiotic stress tolerance in crops: a review. *Plants*, 9(10), 1374.
- Zhao, C., Liu, B., Piao, S., Wang, X., Lobell, D. B., Huang, Y., Huang, M., Yao, Y., Bassu, S., & Ciaia, P. (2017). Temperature increase reduces global yields of major crops in four independent estimates. *Proceedings of the National Academy of sciences*, 114(35), 9326-9331.
- Zörb, C., Ludewig, U., & Hawkesford, M. J. (2018). Perspective on wheat yield and quality with reduced nitrogen supply. *Trends in plant science*, 23(11), 1029-1037.
- Zubair, M., Shakir, M., Ali, Q., Rani, N., Fatima, N., Farooq, S., Shafiq, S., Kanwal, N., Ali, F., & Nasir, I. A. (2016). Rhizobacteria and phytoremediation of heavy metals. *Environmental Technology Reviews*, 5(1), 112-119.

Declaration

Conflict of interest

The researchers affirm that there were no financial or commercial ties that might be seen as a potential conflict of interest throughout the research's execution.

Data Availability statement

All data generated or analyzed during the study have been included in the manuscript.

Ethics approval and consent to participate

These aspects are not applicable in this research.

Consent for publication

Not applicable

Funding

There were no sources providing support, for this research.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, [Creative Commons Attribution-NonCommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/). © The Author(s) 2023